



ACTA1 gene

actin, alpha 1, skeletal muscle

Normal Function

The *ACTA1* gene provides instructions for making a protein called skeletal alpha (α)-actin, which is part of the actin protein family. Actin proteins are important for cell movement and the tensing of muscle fibers (muscle contraction). These proteins also help maintain the cytoskeleton, which is the structural framework that determines cell shape and organizes cell contents.

Skeletal α -actin plays an important role in skeletal muscles, which are muscles that the body uses for movement. Within skeletal muscle cells, skeletal α -actin is an essential component of structures called sarcomeres. Sarcomeres are composed of thin filaments made up of actin and thick filaments made up of another protein called myosin. Attachment (binding) and release of the overlapping thick and thin filaments allows them to move relative to each other so that the muscles can contract.

Health Conditions Related to Genetic Changes

actin-accumulation myopathy

At least nine mutations in the *ACTA1* gene have been identified in people with actin-accumulation myopathy. Most of these mutations change single protein building blocks (amino acids) in the skeletal α -actin protein sequence.

Researchers suggest that *ACTA1* gene mutations that cause actin-accumulation myopathy may affect the way the actin binds to ATP. ATP is a molecule that supplies energy for cells' activities and is important in the formation of thin filaments from individual actin molecules. Dysfunctional actin-ATP binding may result in abnormal thin filament formation and impair muscle contraction, leading to muscle weakness and the other signs and symptoms of actin-accumulation myopathy.

cap myopathy

At least one *ACTA1* gene mutation has been identified as a cause of cap myopathy. The mutation replaces the amino acid methionine with the amino acid valine at position 47 in the protein sequence, written as Met47Val or M47V. The resulting abnormal protein may interfere with the proper assembly of thin filaments. Cap myopathy is characterized by the presence of cap-like structures in muscle cells, and these structures are composed of disorganized thin filaments. The abnormal filament

structure likely impairs the ability of skeletal muscles to contract, resulting in muscle weakness and the other signs and symptoms of cap myopathy.

congenital fiber-type disproportion

At least seven mutations in the *ACTA1* gene have been found to cause congenital fiber-type disproportion, a disorder that causes general muscle weakness that typically does not worsen over time. The mutations that cause this condition change single amino acids in skeletal α -actin. These mutations lead to the production of an abnormal actin protein, which interferes with the function of normal actin proteins in the sarcomere. As a result, the function of the sarcomere is impaired, which disrupts muscle contraction. Inefficient muscle contraction leads to muscle weakness in people with congenital fiber-type disproportion.

intranuclear rod myopathy

At least 13 mutations in the *ACTA1* gene have been identified in people with intranuclear rod myopathy. These mutations change single amino acids in the skeletal α -actin protein sequence.

ACTA1 gene mutations that cause intranuclear rod myopathy result in rod-shaped accumulations of actin in the nucleus of muscle cells. Normally, most actin is found in the fluid surrounding the nucleus (the cytoplasm), with small amounts in the nucleus itself. Researchers suggest that the *ACTA1* gene mutations that cause intranuclear rod myopathy may interfere with the normal transport of actin between the nucleus and the cytoplasm, resulting in the accumulation of actin in the nucleus and the formation of intranuclear rods. Abnormal accumulation of actin in the nucleus of muscle cells and a corresponding reduction of available actin in muscle fibers may impair muscle contraction and lead to the muscle weakness seen in intranuclear rod myopathy.

A few *ACTA1* gene mutations that have been identified in people with intranuclear rod myopathy have also been found in people with actin-accumulation myopathy. It is unclear how the same mutation can cause two different conditions.

nemaline myopathy


More than 170 mutations in the *ACTA1* gene have been found to cause nemaline myopathy. Nemaline myopathy is the most common muscle disorder associated with *ACTA1* gene mutations. Some of the mutations that cause this disorder alter the structure or function of skeletal α -actin, causing the protein to cluster together and form clumps (aggregates). These aggregates interfere with the normal functioning of muscle cells. Other *ACTA1* gene mutations prevent the production of any skeletal α -actin, impairing the muscle cells' ability to contract. *ACTA1* gene mutations that

cause nemaline myopathy impair muscle contraction, causing weakness and the other features of this condition.

Chromosomal Location

Cytogenetic Location: 1q42.13, which is the long (q) arm of chromosome 1 at position 42.13

Molecular Location: base pairs 229,431,245 to 229,434,096 on chromosome 1 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- ACTA
- ACTS_HUMAN
- alpha skeletal muscle actin
- ASMA

Additional Information & Resources

Educational Resources

- Molecular Cell Biology (fourth edition, 2000): Muscle Contraction
<https://www.ncbi.nlm.nih.gov/books/NBK9961/#A1791>

GeneReviews

- Congenital Fiber-Type Disproportion
<https://www.ncbi.nlm.nih.gov/books/NBK1259>
- Nemaline Myopathy
<https://www.ncbi.nlm.nih.gov/books/NBK1288>

Genetic Testing Registry

- GTR: Genetic tests for ACTA1
<https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=58%5Bgeneid%5D>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28ACTA1%5BTI%5D%29+OR+%28alpha+skeletal+muscle+actin%5BTI%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- ACTIN, ALPHA, SKELETAL MUSCLE 1
<http://omim.org/entry/102610>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_ACTA1.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=ACTA1%5Bgene%5D>
- HGNC Gene Family: Actins
<http://www.genenames.org/cgi-bin/genefamilies/set/929>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=129
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/58>
- UniProt
<http://www.uniprot.org/uniprot/P68133>

Sources for This Summary

- OMIM: ACTIN, ALPHA, SKELETAL MUSCLE 1
<http://omim.org/entry/102610>
- Agrawal PB, Strickland CD, Midgett C, Morales A, Newburger DE, Poulos MA, Tomczak KK, Ryan MM, Iannaccone ST, Crawford TO, Laing NG, Beggs AH. Heterogeneity of nemaline myopathy cases with skeletal muscle alpha-actin gene mutations. *Ann Neurol*. 2004 Jul;56(1):86-96.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15236405>
- Castiglioni C, Cassandrini D, Fattori F, Bellacchio E, D'Amico A, Alvarez K, Gejman R, Diaz J, Santorelli FM, Romero NB, Bertini E, Bevilacqua JA. Muscle magnetic resonance imaging and histopathology in ACTA1-related congenital nemaline myopathy. *Muscle Nerve*. 2014 Dec;50(6):1011-6. doi: 10.1002/mus.24353. Epub 2014 Oct 30.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25088345>

- Clarke NF, Ilkovski B, Cooper S, Valova VA, Robinson PJ, Nonaka I, Feng JJ, Marston S, North K. The pathogenesis of ACTA1-related congenital fiber type disproportion. *Ann Neurol*. 2007 Jun; 61(6):552-61.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17387733>
- Feng JJ, Marston S. Genotype-phenotype correlations in ACTA1 mutations that cause congenital myopathies. *Neuromuscul Disord*. 2009 Jan;19(1):6-16. doi: 10.1016/j.nmd.2008.09.005. Epub 2008 Oct 30. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18976909>
- GeneReview: Congenital Fiber-Type Disproportion
<https://www.ncbi.nlm.nih.gov/books/NBK1259>
- GeneReview: Nemaline Myopathy
<https://www.ncbi.nlm.nih.gov/books/NBK1288>
- Hung RM, Yoon G, Hawkins CE, Halliday W, Biggar D, Vajsar J. Cap myopathy caused by a mutation of the skeletal alpha-actin gene ACTA1. *Neuromuscul Disord*. 2010 Apr;20(4):238-40. doi: 10.1016/j.nmd.2010.01.011. Epub 2010 Mar 19. Erratum in: *Neuromuscul Disord*.2010 Aug;20(8): 567.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20303757>
- Ilkovski B, Clement S, Sewry C, North KN, Cooper ST. Defining alpha-skeletal and alpha-cardiac actin expression in human heart and skeletal muscle explains the absence of cardiac involvement in ACTA1 nemaline myopathy. *Neuromuscul Disord*. 2005 Dec;15(12):829-35. Epub 2005 Nov 8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16288873>
- Kaimaktchiev V, Goebel H, Laing N, Narus M, Weeks D, Nixon R. Intranuclear nemaline rod myopathy. *Muscle Nerve*. 2006 Sep;34(3):369-72.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16477620>
- Koy A, Ilkovski B, Laing N, North K, Weis J, Neuen-Jacob E, Mayatepek E, Voit T. Nemaline myopathy with exclusively intranuclear rods and a novel mutation in ACTA1 (Q139H). *Neuropediatrics*. 2007 Dec;38(6):282-6. doi: 10.1055/s-2008-1065356.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18461503>
- Laing NG, Clarke NF, Dye DE, Liyanage K, Walker KR, Kobayashi Y, Shimakawa S, Hagiwara T, Ouvrier R, Sparrow JC, Nishino I, North KN, Nonaka I. Actin mutations are one cause of congenital fibre type disproportion. *Ann Neurol*. 2004 Nov;56(5):689-94.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15468086>
- Laing NG, Dye DE, Wallgren-Pettersson C, Richard G, Monnier N, Lillis S, Winder TL, Lochmüller H, Graziano C, Mitrani-Rosenbaum S, Twomey D, Sparrow JC, Beggs AH, Nowak KJ. Mutations and polymorphisms of the skeletal muscle alpha-actin gene (ACTA1). *Hum Mutat*. 2009 Sep;30(9): 1267-77. doi: 10.1002/humu.21059.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19562689>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2784950/>
- Nowak KJ, Sewry CA, Navarro C, Squier W, Reina C, Ricoy JR, Jayawant SS, Childs AM, Dobbie JA, Appleton RE, Mountford RC, Walker KR, Clement S, Barois A, Muntoni F, Romero NB, Laing NG. Nemaline myopathy caused by absence of alpha-skeletal muscle actin. *Ann Neurol*. 2007 Feb; 61(2):175-84.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17187373>

- Ochala J. Thin filament proteins mutations associated with skeletal myopathies: defective regulation of muscle contraction. *J Mol Med (Berl)*. 2008 Nov;86(11):1197-204. doi: 10.1007/s00109-008-0380-9. Epub 2008 Jun 24. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18574571>
- Schröder JM, Durling H, Laing N. Actin myopathy with nemaline bodies, intranuclear rods, and a heterozygous mutation in ACTA1 (Asp154Asn). *Acta Neuropathol*. 2004 Sep;108(3):250-6. Epub 2004 Jun 24.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15221331>
- Sparrow JC, Nowak KJ, Durling HJ, Beggs AH, Wallgren-Pettersson C, Romero N, Nonaka I, Laing NG. Muscle disease caused by mutations in the skeletal muscle alpha-actin gene (ACTA1). *Neuromuscul Disord*. 2003 Sep;13(7-8):519-31. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12921789>
- Wallefeld W, Krause S, Nowak KJ, Dye D, Horváth R, Molnár Z, Szabó M, Hashimoto K, Reina C, De Carlos J, Rosell J, Cabello A, Navarro C, Nishino I, Lochmüller H, Laing NG. Severe nemaline myopathy caused by mutations of the stop codon of the skeletal muscle alpha actin gene (ACTA1). *Neuromuscul Disord*. 2006 Oct;16(9-10):541-7. Epub 2006 Sep 1.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16945536>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/ACTA1>

Reviewed: May 2016

Published: February 14, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services